

Relationship Between Physical and Biological Properties on the Microscale: A Cross-Comparison Between Differing Coastal Domains

Timothy J. Cowles

College of Oceanic and Atmospheric Sciences

104 Oceanography Admin Bldg

Oregon State University

Corvallis, OR 97331-5503

Office: (541) 737-3966 FAX: (541) 737-2064 email: tjc@coas.oregonstate.edu

Grant # N00014-09-1-0389

Jonathan Nash

College of Oceanic and Atmospheric Sciences

104 Oceanography Admin Bldg

Oregon State University

Corvallis, OR 97331-5503

Office: (541) 737-4573 FAX: (541) 737-2064 email: nash@coas.oregonstate.edu

James Moum

College of Oceanic and Atmospheric Sciences

104 Oceanography Admin Bldg

Oregon State University

Corvallis, OR 97331-5503

Office: (541) 737-2553 FAX: (541) 737-2064 email: moum@coas.oregonstate.edu

LONG-TERM GOALS

Our long-term goal is to quantify the interactions between small-scale biological and physical processes within the upper ocean.

OBJECTIVES

Attenuation of light and sound propagation in the upper ocean is tightly linked to the patterns of vertical distribution of particulate matter (primarily detritus, phytoplankton, and zooplankton). Local maxima in these distributions frequently occur within the euphotic zone, and are often concentrated within depth intervals that are less than 1-2m thick (e.g., Dekshenieks et al. 2001, Cowles 2004). Of particular interest are the steep vertical gradients of stratification and/or shear that often exist at the boundaries of these thin plankton layers, and which are implicated in their formation and maintenance (e.g., Stacey et al. 2007, Birch et al. 2008). We are extending the analysis of thin plankton layers by examining the microstructure datasets obtained over the past decade by the Ocean Mixing Group at Oregon State University. Tens of thousands of profiles have been obtained over the central Oregon shelf (COAST), over the New Jersey shelf (SW06), and around and through the buoyant Columbia River plume (RISE). Plots of cross-shelf property distributions have revealed extensive layers of

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2009		2. REPORT TYPE		3. DATES COVERED 00-00-2009 to 00-00-2009	
4. TITLE AND SUBTITLE Relationship Between Physical and Biological Properties on the Microscale: A Cross-Comparison Between Differing Coastal Domains				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon State University, College of Oceanic and Atmospheric Sciences, 104 Oceanography Admin Bldg, Corvallis, OR, 97331-5503				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

enhanced particulate biomass. Over the Oregon continental shelf in 2001, successive cross-shelf transects consistently revealed filaments of turbidity that extended $O(10\text{km})$. Chameleon profiles obtained off the Washington coast in 2006 also detected layers of enhanced chlorophyll fluorescence that appeared to occur in vertical intervals of relatively low TKE dissipation. In the original processing, the physical dynamics in and around these turbidity and chlorophyll features were not examined. We are now addressing the explicit relationships between these features.

Our specific objectives address the mechanisms that form and maintain steep gradients and thin layers of particulate concentration. Our objectives are to:

- examine existing datasets of Chameleon profiles (upwelling domain over Oregon shelf, CR plume, NJ shelf) for persistent layers of particulate matter (defined by turbidity or chlorophyll);
- quantify the relationship between layer characteristics (vertical gradients, thickness, relationship to pycnocline) and physical descriptors (stratification, shear, TKE dissipation, turbulent mixing);
- quantify the spatial patterns, temporal persistence and coherence of the physical descriptors to assess how these permit both persistent layers and regions void of thin layers;
- compare these relationships as expressed within the upwelling system, buoyant plume, and NJ shelf;
- relate regional layer characteristics to local and regional forcing (wind, current jets, etc);
- examine results in the context of recent theoretical developments about layer formation and maintenance.

APPROACH

We address the project objectives through a multi-stage analysis of the various datasets obtained with Chameleon. We have sorted the dataset of profiles to obtain those with quality-controlled turbidity or chlorophyll fluorescence data. Then subsets of those data have been examined for temporal and/or spatial coherence of identifiable features, particularly the steepness of vertical gradients and local maxima in turbidity or chlorophyll fluorescence. Since no automated methods exist for this type of feature extraction, we are developing and validating layer-referenced gradient detection methods for the Chameleon datasets. We are examining the relationship between vertical gradients of particles with stratification, shear, and TKE dissipation, using depth (or density) of layer maxima as the reference level for analysis. Given the high quality of the shear probe data from the Chameleon datasets, we can resolve TKE dissipation into 0.5 m depth intervals relative to the plankton layer boundaries. As we progress in the analysis, we will apply multi-dimensional ordination to examine the parameter ranges within which we observe persistent layers of particles, and determine if regional differences exist in these relationships.

WORK COMPLETED

We have focused the initial analysis work on the several thousand Chameleon profiles obtained off the Oregon coast during 2001 and 2006 that display obvious planktonic thin layers. We have found wavelet analysis useful in the data extraction process. Using a feature-referenced approach, we are

currently compiling statistics on the relationship between small-scale mixing parameters and plankton features.

RESULTS

We illustrate our results to date with figures that relate plankton layers, as represented by fluorescence, to TKE dissipation (epsilon), stratification (N^2) and density. The data were obtained along a transect just west of the mouth of the Columbia River in May 2006.




Figure 1. Vertical section across a portion of the Columbia River plume in May 2006, illustrating 110 successive Chameleon profiles during an east-west transit. X-AXIS: profile #; Y-AXIS: depth(m). Ship turned around at profile #211. Upper panel shows the fluorescence layer (log units), with maximum value represented by thin black line. Successive panels show the relationship of that fluorescence maximum with TKE dissipation, buoyancy frequency, optical backscatter, and density.




Figure 2. *Compilation of feature-referenced statistics of fluorescence, density, stratification and TKE dissipation from the Chameleon transect illustrated in Figure 1.*

Our initial results suggest the layer resides in a transitional region below the stratified, turbulent plume of the Columbia River. The plankton layer is capped above by relatively higher stratification, while dissipation and mixing are considerably lower below the layer. We suspect that strain and dissipation play competing roles in layer evolution.

IMPACT/APPLICATION

We expect that this work will reinforce the need for coincident measurements of microstructure physics and biology.

TRANSITIONS

None at this time.

RELATED PROJECTS

REFERENCES

Birch, D.A., W.R. Young and P.J.S. Franks. 2008. Thin layers of plankton: Formation by shear and death by diffusion. *Deep-Sea Res.* 55: 277-295

- Cowles, T.J. 2004. Planktonic layers: physical and biological interactions on the small scale. In: *Handbook of Scaling Methods in Aquatic Ecology: Measurements, Analysis, Simulation*. CRC Press, Boca Raton, FL. , pp 31-49.
- Dekshenieks, M.M., Donaghay, P.L., Sullivan, J.M., Rines, J.E.B., Osborn, T.R., and Twardowski, M.S. 2001. Temporal and spatial occurrence of thin phytoplankton layers in relation to physical processes. *Mar. Ecol. Prog. Ser.*, 223: 61
- Stacey, M.T., M.A. McManus, and J.V. Steinbuck. 2007. Convergences and divergences and thin layer formation and maintenance. *Limnol. Oceanogr.*, 52: 1523–1532

PUBLICATIONS (refereed)

None